

Extended Quark Potential Model from Random Phase Approximation

Weizhen Deng, Xiaolin Chen, Dahai Lu, and Liming Yang

Department of Physics, Peking University, Beijing 100871, China

Abstract

The quark potential model is extended to include the sea quark excitation using the random phase approximation (RPA). The effective quark interaction preserves the important Quantum Chromodynamics (QCD) properties — chiral symmetry and confinement simultaneously. A primary qualitative analysis shows that the π meson as a well-known typical Goldstone boson and the other mesons made up of valence $q\bar{q}$ quark pair such as the ρ meson can also be described in this extended quark potential model.

PACS numbers: 12.39.Pn, 14.40.Aq

I. INTRODUCTION

The Quantum Chromodynamics (QCD) is widely believed to be the correct theory of strong interaction nowadays. Due to the complication of non-Abelian gauge theory, in the studies of hadron structure, models are often adopted to approximate the strong interaction. Two kinds of models are often used, each incorporates several important QCD features. The quark potential model based on the confinement quark picture has been impressively successful in hadron spectra [1], except for the π meson. Recently, it was shown that for the π meson, which is a well-known Goldstone boson, the explanation of quark potential model, i.e. mesons merely made up of $q\bar{q}$ pairs of valence quarks, is not well satisfied and the important sea quark contribution must be considered [2]. On the other hand, the Nambu–Jona-Lasinio (NJL) model [3] and several of its extensions [4, 6] can well describe the π meson as a Goldstone boson which is a conclusion of QCD chiral symmetry. The sea quark contributions to the meson structure are included in the NJL model by using the Bethe-Salpeter equation or its equivalence — Random Phase Approximation (RPA). Also ground states of other mesons such as ρ which are made up of simple valence $q\bar{q}$ quark pair can be described by the extensions to the original NJL model. However, the NJL model lacks the important QCD property of quark confinement, so the excitation states of mesons cannot be accounted in the model, such as π' , ρ' , ρ'' , etc.

In this paper, we try to incorporate the important QCD chiral symmetry into the quark potential model by comprising the sea quark contribution. To incorporate the chiral symmetry, we start from an effective quark hamiltonian taken from extended NJL models which is chiral invariant. To consider the sea quark contribution, we adopt the RPA approximation. This derives a hamiltonian with two channels, i.e. the ordinary valence quark channel and a new sea quark channel, which is an extension to the quark potential model. This extended quark potential model with RPA approximation demands a new potential which couples the valence quark channel with the sea quark channel we call coupling potential. Then a preliminary numerical analysis is made to illustrate that the Goldstone π meson and also ρ meson can be well described together in the extended quark potential model.

In the next section, We obtain the extended quark potential model — the RPA approximation is used to derive the two channel hamiltonian and the so called coupling potential. In sec. 3, a preliminary analysis is made to the π and ρ mesons. Finally, some brief summary

and discussion are given.

II. EXTENDED QUARK POTENTIAL MODEL WITH RPA

In this section, we will incorporate the QCD chiral symmetry into the original quark potential model. To avoid the complication of QCD, our start point will be an effective quark interaction. A good choice of the effective quark interaction with chiral symmetry is of the NJL type. However, despite of much success of the original NJL model, its contact quark interaction is too simplified as an effective quark interaction which does not possess the important QCD feature of color confinement. The original NJL model has been modified by several authors to include the nonlocal effective quark interaction [4, 6].

Consider a quark nonlocal action where the QCD strong interaction is approximated by the quark effective interaction after the complicated degrees of freedom of gluon are eliminated as in the extended NJL models. Its general form is

$$A = \int d^4x \bar{\Psi}(x)(\gamma \cdot p - m_0)\Psi(x) + \frac{1}{2} \sum_{\epsilon} \int d^4x d^4y K_{\epsilon}(x-y) \bar{\Psi}(x) \Gamma_{\epsilon} \Psi(x) \bar{\Psi}(y) \Gamma_{\epsilon} \Psi(y), \quad (1)$$

where $K_{\epsilon}(x)$ are the interaction kernel functions and Γ_{ϵ} are tensor products of Dirac, flavor (in this paper, we will restrict to iso-spin $SU(2)$ symmetry only) and color ($SU(3)$) matrices which represent the quark interaction channels. The quark interaction in this quark action will be chosen to have chiral symmetry and with quark confinement in the next section.

As the action is nonlocal, the exact hamiltonian will be very complicated and also time dependent. In the present paper we shall only consider the non-relativistic limit. We can adopt a simple, time independent hamiltonian with an instantaneous approximation. After approximating the kernel function as

$$K_{\epsilon}(x-y) \approx K_{\epsilon}(\mathbf{x} - \mathbf{y}) \delta(x^0 - y^0), \quad (2)$$

the model hamiltonian becomes

$$H = H_2 + H_4, \quad (3)$$

$$H_2 = \int d^3x \Psi^{\dagger}(\mathbf{x})(-i\boldsymbol{\alpha} \cdot \boldsymbol{\nabla} + \beta m_q)\Psi(\mathbf{x}), \quad (4)$$

$$H_4 = -\frac{1}{2} \sum_{\epsilon} \int d^3x d^3y K_{\epsilon}(\mathbf{x} - \mathbf{y}) \bar{\Psi}(\mathbf{x}) \Gamma_{\epsilon} \Psi(\mathbf{x}) \bar{\Psi}(\mathbf{y}) \Gamma_{\epsilon} \Psi(\mathbf{y}), \quad (5)$$

where $\Psi(\mathbf{x}) \equiv \Psi(\mathbf{x}, 0)$ which can be expressed in quark creation and annihilation operators

$$\Psi(\mathbf{x}) = \sum_s \int \frac{d^3k}{(2\pi)^3} \frac{1}{2k^0} \left[b(\vec{k}, s) u(\vec{k}, s) e^{i\mathbf{k} \cdot \mathbf{x}} + d^\dagger(\vec{k}, s) v(\vec{k}, s) e^{-i\mathbf{k} \cdot \mathbf{x}} \right]. \quad (6)$$

The hamiltonian from the instantaneous approximation breaks the important Lorentz invariance. We shall only use it to study the hadron structure in rest frame. For a moving hadron, however, we shall use the Lorentz boost technique to restore the Lorentz invariance in future study.

Apart from a constant, the hamiltonian (3) can be written in normal order form as

$$H = H_0 + : H_4 : \quad (7)$$

where H_0 represents the single particle energy of constituent quark and it can be re-written formly as

$$H_0 = \int d^3x \Psi^\dagger(\mathbf{x}) \left[-iA(-\nabla^2) \boldsymbol{\alpha} \cdot \nabla + \beta B(-\nabla^2) \right] \Psi(\mathbf{x}). \quad (8)$$

The function forms of A and B operators should be obtained from the Mean Field Approximation (MFA) and their momentum dependence are due to the Lorentz invariance breaking from instantaneous approximation. Here for simplification we shall take the A and B operators as constants. This only introduce one parameter of the constituent quark mass m_c . The constituent quark single energy is simplified as

$$H_0 = \int d^3x \Psi^\dagger(\mathbf{x}) [-i\boldsymbol{\alpha} \cdot \nabla + \beta m_c] \Psi(\mathbf{x}). \quad (9)$$

It is generally accepted that hadrons are excited states of the QCD vacuum. To account the sea quark contribution, we consider a meson as a superposition of creation and annihilation of $q\bar{q}$ pairs on the vacuum. Thus the meson states are treated as an excitation modes of the vacuum of the RPA type:

$$|Q\rangle = Q^\dagger |0\rangle, \quad (10)$$

where $Q^\dagger$ is the excitation operator. For a meson in the rest frame, $Q^\dagger$ can be expressed in quark and antiquark creation and annihilation operators

$$Q^\dagger = \sum_{s_1 s_2} \int \frac{d^3k}{(2\pi)^3} \frac{1}{2k^0} \left[x(\mathbf{k}, s_1, s_2) b^\dagger(\mathbf{k} s_1) d^\dagger(-\mathbf{k} s_2) \right. \\ \left. + y(\mathbf{k}, s_1, s_2) \tilde{d}(-\mathbf{k} s_1) \tilde{b}(\mathbf{k} s_2) \right], \quad (11)$$

where x and y are the x-amplitude and y-amplitude of the RPA wave function respectively and the time reversal operators are

$$\tilde{b}(\mathbf{k}s) \equiv (-1)^{1/2-s}b(\mathbf{k}-s), \quad (12)$$

$$\tilde{d}(\mathbf{k}s) \equiv (-1)^{1/2-s}d(\mathbf{k}-s). \quad (13)$$

The x- and y-amplitudes can be solved from the RPA equation of motion

$$\langle 0 | [\delta Q, [H, Q^\dagger]] | 0 \rangle = E \langle 0 | [\delta Q, Q^\dagger] | 0 \rangle. \quad (14)$$

After inserting the hamiltonian eq. (7) into the RPA equation of motion, we obtain the equations for the x- and y- amplitudes

$$\begin{aligned} & (E - 2k^0) x(\mathbf{k}, s_1, s_2) \\ &= - \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(0) \\ & \quad \left\{ \bar{u}(\mathbf{k}s_1) \Gamma_{\epsilon} v(-\mathbf{k}s_2) \bar{v}(-\mathbf{k}'s_4) \Gamma_{\epsilon} u(\mathbf{k}'s_3) x(\mathbf{k}', s_3, s_4) \right. \\ & \quad \left. - \bar{u}(\mathbf{k}s_1) \Gamma_{\epsilon} v(-\mathbf{k}s_2) \tilde{u}(\mathbf{k}'s_4) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}'s_3) y(\mathbf{k}', s_3, s_4) \right\} \\ &+ \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\ & \quad \left\{ \bar{u}(\mathbf{k}s_1) \Gamma_{\epsilon} u(\mathbf{k}'s_3) \bar{v}(-\mathbf{k}'s_4) \Gamma_{\epsilon} v(-\mathbf{k}s_2) x(\mathbf{k}', s_3, s_4) \right. \\ & \quad \left. - \bar{u}(\mathbf{k}s_1) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}'s_3) \tilde{u}(\mathbf{k}'s_4) \Gamma_{\epsilon} v(-\mathbf{k}s_2) y(\mathbf{k}', s_3, s_4) \right\}, \end{aligned} \quad (15)$$

and

$$\begin{aligned} & (E + 2k^0) y(\mathbf{k}, s_1, s_2) \\ &= - \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(0) \\ & \quad \left\{ \tilde{v}(-\mathbf{k}s_1) \Gamma_{\epsilon} \tilde{u}(\mathbf{k}s_2) \bar{v}(-\mathbf{k}'s_4) \Gamma_{\epsilon} u(\mathbf{k}'s_3) x(\mathbf{k}', s_3, s_4) \right. \\ & \quad \left. - \tilde{v}(-\mathbf{k}s_1) \Gamma_{\epsilon} \tilde{u}(\mathbf{k}s_2) \tilde{u}(\mathbf{k}'s_4) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}'s_3) y(\mathbf{k}', s_3, s_4) \right\} \\ &+ \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\ & \quad \left\{ \tilde{v}(-\mathbf{k}s_1) \Gamma_{\epsilon} u(\mathbf{k}'s_3) \bar{v}(-\mathbf{k}'s_4) \Gamma_{\epsilon} \tilde{u}(\mathbf{k}s_2) x(\mathbf{k}', s_3, s_4) \right. \\ & \quad \left. - \tilde{v}(-\mathbf{k}s_1) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}'s_3) \tilde{u}(\mathbf{k}'s_4) \Gamma_{\epsilon} \tilde{u}(\mathbf{k}s_2) y(\mathbf{k}', s_3, s_4) \right\}, \end{aligned} \quad (16)$$

where $\tilde{u}$ and $\tilde{v}$ are the time reversal of spinors u and v respectively. In each equation, the first summation (direct term) can be absorbed into the second summation (exchange term)

by Fierz rearrangement, then equations briefly become

$$\begin{aligned}
& (E - 2k^0) x(\mathbf{k}, s_1, s_2) \\
&= \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\
& \quad \{ \bar{u}(\mathbf{k} s_1) \Gamma_{\epsilon} u(\mathbf{k}' s_3) \bar{v}(-\mathbf{k}' s_4) \Gamma_{\epsilon} v(-\mathbf{k} s_2) x(\mathbf{k}', s_3, s_4) \\
& \quad - \bar{u}(\mathbf{k} s_1) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}' s_3) \tilde{u}(\mathbf{k}' s_4) \Gamma_{\epsilon} v(-\mathbf{k} s_2) y(\mathbf{k}', s_3, s_4) \}, \tag{17}
\end{aligned}$$

and

$$\begin{aligned}
& (E + 2k^0) y(\mathbf{k}, s_1, s_2) \\
&= \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\
& \quad \{ \tilde{\bar{v}}(-\mathbf{k} s_1) \Gamma_{\epsilon} u(\mathbf{k}' s_3) \bar{v}(-\mathbf{k}' s_4) \Gamma_{\epsilon} \tilde{u}(\mathbf{k} s_2) x(\mathbf{k}', s_3, s_4) \\
& \quad - \tilde{\bar{v}}(-\mathbf{k} s_1) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}' s_3) \tilde{u}(\mathbf{k}' s_4) \Gamma_{\epsilon} \tilde{u}(\mathbf{k} s_2) y(\mathbf{k}', s_3, s_4) \}. \tag{18}
\end{aligned}$$

Now We introduce two effective potentials U and V , such that the matrix elements are

$$\begin{aligned}
\langle \mathbf{k}, s_1, s_2 | U | \mathbf{k}', s_3, s_4 \rangle &\equiv \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\
& \quad \bar{u}(\mathbf{k} s_1) \Gamma_{\epsilon} u(\mathbf{k}' s_3) \bar{v}(-\mathbf{k}' s_4) \Gamma_{\epsilon} v(-\mathbf{k} s_2), \tag{19}
\end{aligned}$$

$$\begin{aligned}
\langle \mathbf{k}, s_1, s_2 | V | \mathbf{k}', s_3, s_4 \rangle &\equiv \frac{1}{2k^0} \frac{1}{2k'^0} \sum_{\epsilon} K_{\epsilon}(\mathbf{k} - \mathbf{k}') \\
& \quad \bar{u}(\mathbf{k} s_1) \Gamma_{\epsilon} \tilde{v}(-\mathbf{k}' s_3) \tilde{u}(\mathbf{k}' s_4) \Gamma_{\epsilon} v(-\mathbf{k} s_2). \tag{20}
\end{aligned}$$

From the basic symmetries of parity, charge conjugation and time reversal of quark interaction, and the definition eqs. (19) and (20), one can obtain the hermicity of effective potentials U and V

$$U^{\dagger} = U, \tag{21}$$

$$V^{\dagger} = V, \tag{22}$$

and also

$$\begin{aligned}
& \langle \mathbf{k}, s_1, s_2 | U | \mathbf{k}', s_3, s_4 \rangle^* \\
&= (-1)^{1/2-s_1} (-1)^{1/2-s_2} (-1)^{1/2-s_3} (-1)^{1/2-s_4} \\
& \quad \langle \mathbf{k}, -s_2, -s_1 | U | \mathbf{k}', -s_4, -s_3 \rangle. \tag{23}
\end{aligned}$$

Finally, the RPA equations are written as

$$(E - 2k^0) x(\mathbf{k}, s_1, s_2)$$

$$\begin{aligned}
&= \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} [\langle \mathbf{k}, s_1, s_2 | U | \mathbf{k}', s_3, s_4 \rangle x(\mathbf{k}', s_3, s_4) \\
&\quad - \langle \mathbf{k}, s_1, s_2 | V | \mathbf{k}', s_3, s_4 \rangle y(\mathbf{k}', s_3, s_4)],
\end{aligned} \tag{24}$$

and

$$\begin{aligned}
&(E + 2k^0) Y(\mathbf{k}, s_1, s_2) \\
&= \sum_{s_3 s_4} \int \frac{d^3 k'}{(2\pi)^3} [\langle \mathbf{k}, s_1, s_2 | V | \mathbf{k}', s_3, s_4 \rangle x(\mathbf{k}', s_3, s_4) \\
&\quad - \langle \mathbf{k}, s_1, s_2 | U | \mathbf{k}', s_3, s_4 \rangle y(\mathbf{k}', s_3, s_4)].
\end{aligned} \tag{25}$$

In the above RPA equations, the potential U interacts between the same RPA amplitudes (x-amplitude in eq. (24) and y-amplitude in eq. (25)). The x-amplitude represents the contribution of valence quarks, that is just the wave-function of valence quarks in the potential model. We conclude that the U potential is just the quark potential in the potential model. On the other hand, potential V interacts between the different RPA amplitudes. This is a new potential and we call it the coupling potential.

The above RPA approximation may be considered as an extension to the potential model. The extended potential model includes two channels — the ordinary valence quark channel and an extended sea quark channel. The valence quark channel which represents the contribution of valence quarks is described by the x-amplitudes. The sea quark channel representing the contribution of sea quarks is described by the y-amplitudes. The ordinary quark potential U interacts within both individual channel. The new coupling potential V couples the two channels together.

The potentials U and V will be evaluated from the kernels in action (1). Following ref. [7], from the spinor structure of Lorentz invariant, the kernels can be divided into five types, i.e., scalar (S), pseudo-scalar (P), vector (V), axial-vector (A), and tensor (T). We write

$$K = S1 \otimes 1 + P i\gamma^5 \otimes i\gamma^5 + V \gamma^\mu \otimes \gamma_\mu + A \gamma^\mu \gamma^5 \otimes \gamma_\mu \gamma^5 + \frac{1}{2} T \sigma_{\mu\nu} \otimes \sigma^{\mu\nu}. \tag{26}$$

In the non-relativistic limit $\mathbf{k}, \mathbf{k}' \rightarrow 0$, only the following terms survive in the potentials

$$U = -S + V + A \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 + T \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2, \tag{27}$$

$$V = P - V \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 - A - T \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2. \tag{28}$$

where $\boldsymbol{\sigma}$'s are the pauli spin matrices.

III. PRIMARY APPLICATION

In addition to the Lorentz invariance of quark interaction kernels, in the limit of vanishing current quark masses, the kernels must also have the chiral symmetry of QCD. The general quark interaction which is compatible with the chiral symmetry can be found in ref. [5]. Here we simply choose the form of effective quark interaction H_4 as a vector interaction like the one gluon exchange interaction

$$H_4 = \frac{1}{2} \int d^3x d^3y K(\mathbf{x} - \mathbf{y}) \bar{\Psi}(\mathbf{x}) \gamma^\mu \boldsymbol{\lambda}^c \Psi(\mathbf{x}) \bar{\Psi}(\mathbf{y}) \gamma_\mu \boldsymbol{\lambda}^c \Psi(\mathbf{y}). \quad (29)$$

Following the quark potential model [1], the effective quark interaction kernel K may be chosen as a confinement interaction plus the one gluon exchange interaction

$$K(r) = -\frac{3}{16}br + \frac{1}{4} \frac{\alpha_s}{r}. \quad (30)$$

Our choice of confinement interaction differs from ref. [1] in that ours is a vector $\gamma^\mu \otimes \gamma_\mu$ interaction. This is because from the effective interaction kernel eq. (26), according to ref. [7], the effective qq interaction is

$$U = S + V - A \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 + T \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2. \quad (31)$$

Notice that scalar interaction $1 \otimes 1$ changes sign from the $q\bar{q}$ (eq. (27)) to qq (eq. (31)). This seems to us that one can not get the confinement for both mesons and baryons simultaneously from a unified interaction kernel.

To actually perform calculations with these potentials, we would like to diagonalize the hamiltonian matrix in the harmonic-oscillator basis following ref. [1]. However, in this primary analysis, we simply approximate the meson ground states using the ground state wave function of harmonic-oscillator

$$x(\mathbf{k}, s_1, s_2) = X \Psi_{000}(k) \langle 1/2 s_1 1/2 s_2 | SM_S \rangle, \quad (32)$$

$$y(\mathbf{k}, s_1, s_2) = Y \Psi_{000}(k) \langle 1/2 s_1 1/2 s_2 | SM_S \rangle, \quad (33)$$

where SM_S is the spin of the meson, and

$$\Psi_{000}(k) = \frac{1}{(\sqrt{\pi}\beta)^{3/2}} e^{-k^2/2\beta^2}. \quad (34)$$

From the above simplification and the effective hamiltonian eq. (29), the RPA equations reduce to

$$\begin{bmatrix} A & B \\ -B & -A \end{bmatrix} \begin{bmatrix} X \\ Y \end{bmatrix} = m \begin{bmatrix} X \\ Y \end{bmatrix}, \quad (35)$$

where

$$A = 2\langle k^0 \rangle + \langle K \rangle, \quad (36)$$

$$B = -\langle K \rangle \langle \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 \rangle, \quad (37)$$

and

$$\langle \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 \rangle = 2S(S+1) - 3. \quad (38)$$

In above formula, $\langle \dots \rangle$ takes the average value. In nonrelativistic limit $\mathbf{k} \rightarrow 0$, hence $k^0 = \sqrt{\mathbf{k}^2 + m_c^2} \approx m_c$, so we have approximately

$$A \approx 2m_c + \langle K \rangle. \quad (39)$$

By introducing

$$\alpha = X + Y, \quad (40)$$

$$\beta = X - Y, \quad (41)$$

the RPA equations become

$$(A + B)\alpha = m\beta, \quad (42)$$

$$(A - B)\beta = m\alpha. \quad (43)$$

For a preliminary analysis, we will only consider two typical mesons — the well-known Goldstone boson π and the vector meson ρ which is well described in the original quark potential model as $q\bar{q}$ valence quark pair. For π , its orbital angular momentum $L = 0$, spin $S = 0$, we have

$$2(m_c + 2\langle K \rangle)\alpha_\pi = m_\pi\beta_\pi, \quad (44)$$

$$2(m_c - \langle K \rangle)\beta_\pi = m_\pi\alpha_\pi. \quad (45)$$

The chiral symmetry gives a relation between constituent quark mass m_c and average value of effective quark interaction $\langle K \rangle$. From MFA

$$m_c = m_q + \Sigma, \quad (46)$$

where m_q is the current quark mass, Σ is the Hatree-Fock quark self-energy. In the chiral limit $m_q \rightarrow 0$, the pion will be an exact massless Goldstone boson, i.e., $m_\pi \rightarrow 0$. From the equations of pion, eqs. (44) and (45), we obtain in the chiral limit where $m_q = 0$, $m_\pi = 0$,

$$\Sigma = \langle K \rangle. \quad (47)$$

Finally

$$m_c = m_q + \langle K \rangle. \quad (48)$$

Back to equations of π meson, we obtain

$$m_\pi = 2\sqrt{m_q(3m_c - 2m_q)}, \quad (49)$$

$$\frac{\alpha_\pi}{\beta_\pi} = \sqrt{\frac{m_q}{3m_c - 2m_q}}. \quad (50)$$

For ρ meson, its $L = 0$ and $S = 1$, then

$$2m_c\alpha_\rho = m_\rho\beta_\rho, \quad (51)$$

$$2(m_c + \langle K \rangle)\beta_\rho = m_\rho\alpha_\rho. \quad (52)$$

We obtain

$$m_\rho = 2\sqrt{m_c(2m_c - m_q)}, \quad (53)$$

$$\frac{\alpha_\rho}{\beta_\rho} = \sqrt{\frac{2m_c - m_q}{m_c}}. \quad (54)$$

In the table I, we give two sets of numerical calculations. In the first calculation (*A*), the quark masses (m_q and m_c) are taken to be general accepted values. In the second calculation (*B*), the quark masses are chosen to better fit the experimental data of meson masses (m_π and m_ρ). We believe that both sets of results are reasonable as a primary calculation. The amplitude ratio $Y_{\pi,\rho}/X_{\pi,\rho}$ determines the sea quark contribution vs. the valence quark contribution [9]. Since $X_\pi \approx Y_\pi$ the sea quark contribution is essential to π meson. For ρ meson, $X_\rho \gg Y_\rho$, the sea quark contribution is small.

TABLE I: Numerical calculations (all mass units are MeV).

	m_q	m_c	m_π	m_ρ	X_π/Y_π	X_ρ/Y_ρ
A	5.5	300	140	845	-1.2	5.9
B	6	274	139	771	-1.2	5.9

Since $m_\pi \gg m_q$, we have approximately

$$m_\pi = 2\sqrt{3}\sqrt{m_q m_c}, \quad (55)$$

$$\frac{\alpha_\pi}{\beta_\pi} = \sqrt{\frac{m_q}{3m_c}}, \quad (56)$$

$$m_\rho = 2\sqrt{2}m_c, \quad (57)$$

$$\frac{\alpha_\rho}{\beta_\rho} = \sqrt{2}. \quad (58)$$

Then we find a mass relation

$$m_\pi^2 = \sqrt{18}m_q m_\rho \quad (59)$$

The experimental data $m_\pi = 139\text{MeV}$, $m_\rho = 770\text{MeV}$ combining with the general accepted estimation $m_q = 5.5\text{MeV}$ give

$$\frac{m_\pi^2}{m_q m_\rho} = 4.6, \quad (60)$$

while $\sqrt{18} = 4.2$.

IV. SUMMARY AND DISCUSSION

To take into account of the sea quark contribution to mesons more properly, the original quark potential model is extended which now includes a new sea quark channel similar to the RPA approximation. A new coupling potential is needed to couple the original valence quark channel and the sea quark channel. Through an effective quark interaction which is also chiral invariant as demanded by QCD, the new potential is related to the conventional quark potential in original quark potential model. The constituent quark mass is related to the quark potential also from the chiral symmetry. The primary calculation connected the masses of mesons to masses of quarks (constituent and valence). The sea quark contribution is important to the π meson which is a well-known Goldstone boson. Other mesons, such as

vector ρ meson, where the valence quark contribution is dominant, can also be well described by the same extended quark potential model.

Finally, we shall indicate that in this primary analysis, under the non-relativistic approximation, many important features of quark interaction such as the spin-orbit interaction, the spin-spin interaction and tensor interaction are neglected. Those must be considered in the complete calculation of meson spectroscopy in the future.

Acknowledgments

This work is supported in part by China National Nature Science Foundation.

-
- [1] Stephen Godfrey and Nathan Isgur, Phys. Rev. **D32** (1985) 189.
 - [2] Felipe J. Llanes-Estrada and Stephen R. Cotanch, Phys. Rev. Lett. **84** (2000) 1102.
 - [3] Y. Nambu and G. Jona-Lasinio, Phys. Rev. **122** (1961) 246.
 - [4] V. Bernard, R. Brockmann, M. Schaden, W. Weise and E. Werner, Nucl. Phys. **A412** (1984) 349; V. Bernard, R. Brockmann and W. Weise, Nucl. Phys. **A440** (1985) 605.
 - [5] S. Klimt, M. Lutz, U. Vogl and W. Weise, Nucl. Phys. **A516** (1990) 429.
 - [6] K. Langfeld, C. Kettner and H. Reinhardt, Nucl. Phys. **A608** (1996) 331.
 - [7] D. Gromes, Nucl. Phys. **B131** (1977) 80.
 - [8] Liming Yang, to be published.
 - [9] Weizhen Deng, Xiaolin Chen, Dahai Lu and Liming Yang, Phys. Lett. **B506** (2001) 85.